

$K^\pm \rightarrow \pi^\mp \mu^\pm \mu^\pm$ and doubly-charged Higgs

Charles Picciotto

Department of Physics and Astronomy, University of Victoria, Victoria,

British Columbia, Canada V8W 3P6

(February 1997)

The rate for the lepton-number-violating decay $K^\pm \rightarrow \pi^\mp \mu^\pm \mu^\pm$ is calculated in a model which incorporates doubly-charged Higgs bosons. We find that for reasonable values of the parameters the decay branching ratio may be as large as 2×10^{-16} . Although this is a discouragingly small number, it is of the same order of magnitude as the rate mediated by massive Majorana neutrinos.

I. INTRODUCTION

In the past couple of decades a large amount of effort has been dedicated to the study of double beta decay in nuclei. The motivation has been primarily two-fold: The experimental observation and analysis of the two-neutrino process $(A, Z) \rightarrow (A, Z + 2) + 2e^- + 2\bar{\nu}$ has shed light on the structure of the nuclear matrix elements involved, and the search for the neutrinoless process $(A, Z) \rightarrow (A, Z + 2) + 2e^-$ has provided a powerful tool for the study of the Majorana properties of neutrinos, with considerable speculation about what might be inferred from the actual observation of this lepton-violating decay.

The studies of neutrinoless double beta decay have mostly relied on the assumption that the process would be mediated by massive Majorana neutrinos, with the modern picture being described by leptons which appear in left- and right-handed doublets, the large mass of the right-handed neutrino being responsible for the small left-handed neutrino mass via the seesaw mechanism. In this scheme neutrino double beta decay would proceed as a second-order process, described by the same Hamiltonian that is responsible for the standard weak

processes.

However, the possibility that double-beta decay might be mediated by a different lepton-violating interaction was also considered early on [1], with the suggestion that a $\Delta_{3/2,3/2}$ resonance inside the nucleus might induce a lepton-violating process. Later, a mechanism for $|\Delta L| = 2$ was proposed where the electrons might couple to quarks via a doubly-charged Higgs [2], but subsequent arguments showed that in this case the effective coupling is negligibly small [3]. In fact, the conclusion for nuclear double beta decay was that there are no important $|\Delta L = 2|$ contributions not proportional to m_ν in a $SU(2) \times U(1)$ model with a Higgs triplet.

The Majorana nature of neutrinos can in principle also be tested in the double beta decay of strange mesons, although one would expect very small branching ratios. Experimental searches for nuclear double beta decay put stringent limits on $K^\pm \rightarrow \pi^\mp e^\pm e^\pm$. Calculations for $K^\pm \rightarrow \pi^\mp \mu^\pm \mu^\pm$ have been performed in a model where lepton-number violation is transmitted through the propagation of a heavy Majorana lepton [4], with resulting rates of the order $10^{-16} \text{ sec}^{-1}$. A computation has also been done using a relativistic quark model for mesons [5], which made it possible to consider small neutrino masses. In that case, the resulting rates are several orders of magnitude smaller.

Here we wish to look at the possible contribution to $K^\pm \rightarrow \pi^\mp \mu^\pm \mu^\pm$ that doubly charged Higgs bosons might make, and see how they compare to the previous calculations just mentioned. These doubly-charged Higgs bosons are present in exotic Higgs representations and can have lepton-number-violating couplings to any pair of leptons. They appear in many models, and to provide some focus we shall concentrate on the phenomenology of a model with one $|Y| = 0$ and one $|Y| = 2$ triplet [6]. The features of this model that are relevant for our analysis are the following:

The lepton-number-violating coupling to left-handed leptons is specified by the following Lagrangian:

$$L = ih_{ij}\psi_{iL}^T C\tau_2\Delta\psi_{jL} + h.c. \quad (1)$$

where ψ_{iL} is the two-component leptonic doublet and the Δ is the 2×2 representation of the $Y = 2$ complex triplet. These couplings lead directly to Majorana masses for the neutrinos, so this model does not rely on right-handed lepton partners in order to generate neutrino masses. The Lagrangian in eq. (1) implies a Majorana mass term for the neutrino given by

$$m_{ij} = \frac{h_{ij}s_H v}{\sqrt{2}} \quad (2)$$

where the subscript ij refers to the lepton families, v is related to the neutral Higgs vacuum expectation values, and s_H is the sine of a doublet-triplet mixing angle.

For the case of the electron neutrino, the experimental observation of nuclear double beta decay puts severe limits on the value of m_{ν_e} and as a result on h_{ee} . One obtains [6]

$$s_H h_{ee} \lesssim 6 \times 10^{-12}. \quad (3)$$

Clearly for the electron-electron case we would not expect the model to predict any significant phenomenological consequences. However, in general for some models [7] there are certain niceties that follow from the assumption that the expectation value of the neutral Higgs vanishes. In that case the limit on m_{ν_e} is not relevant, and from Bhabba scattering one may derive the limit [6] [7] [9]

$$h_{ee}^2 < 10^{-5} GeV^{-2} \times m_{\Delta^{++}}^2. \quad (4)$$

Constraints on $h_{\mu\mu}$ can be obtained from the experimental limit on muonium to antimuonium conversion, with the result

$$h_{ee} h_{\mu\mu} < 6 \times 10^{-5} GeV^{-2} \times m_{\Delta^{++}}^2. \quad (5)$$

The physical states can be classified according to their transformation properties, and one finds a fiveplet H_5 (which includes the doubly-charged components), a threeplet H_3 , and two singlets. There are a variety of couplings of the Higgs bosons to fermions and to gauge vector bosons. The ones in which we are potentially interested are

$$H_5^{++} W^- W^- \sim i\sqrt{2} g m_W s_H g_{\mu\nu}$$

$$H_5^{++}H_3^-W^- \sim -i\sqrt{2}c_H e(p_{++} - p_-)_\mu \quad (6)$$

$$H_3^+ q' \bar{q} \sim \frac{g^{s_H}}{2\sqrt{2}m_W c_H} [m_{q'}(1 + \gamma_5) - m_q(1 - \gamma_5)].$$

Finally, we remark that in the limit $s_H \rightarrow 0$ the standard model is regained, with the singlet H_1^0 playing the role of the standard Higgs boson with the standard couplings to fermions and vector mesons.

II. $K^+ \rightarrow \pi^- \mu^+ \mu^+$

The quark-level diagrams that contribute to the decay amplitude are shown in Fig. 1. For this particular model only the phenomenology of left-handed fields was developed [6], but in any case limits on right-handed currents would make them irrelevant here. We adopt the usual form for the hadronic charged weak current

$$\begin{aligned} J^\alpha = & \cos \theta_c [\bar{u}_u \gamma^\alpha (1 - \gamma_5) u_d + \bar{u}_c \gamma^\alpha (1 - \gamma_5) u_s] \\ & + \sin \theta_c [\bar{u}_u \gamma^\alpha (1 - \gamma_5) u_s + \bar{u}_c \gamma^\alpha (1 - \gamma_5) u_d] \end{aligned} \quad (7)$$

The amplitude for Fig. 1(a) is then given by

$$\begin{aligned} T_{1a} = & \frac{ig}{2\sqrt{2}} \langle \pi | J^\mu | 0 \rangle \frac{i}{m_W^2} i\sqrt{2} g m_W s_H \frac{i}{m_{\Delta^{++}}^2} \\ & \times h_{\mu\mu} \langle leptons \rangle \frac{i}{m_W^2} \frac{ig}{2\sqrt{2}} \langle 0 | J^\mu | k \rangle \end{aligned} \quad (8)$$

where $\langle leptons \rangle = \bar{u}(l_1)(1 + \gamma_5)u^c(l_2) - (l_1 \leftrightarrow l_2)$. As usual, the current matrix elements can be expressed as $\langle 0 | J^\mu | k \rangle = i f_k p_k$ and $\langle \pi | J^\mu | 0 \rangle = i f_\pi p_\pi$, where f_π and f_k are the meson decay constants. The amplitude then reduces to

$$T_{1a} = 0.6 G_F^{3/2} m_k f_\pi f_k E_\pi \langle leptons \rangle \frac{s_H h_{\mu\mu}}{m_{\Delta^{++}}^2}. \quad (9)$$

The amplitude for Fig. 1(b) cannot be written down in a similar simple form. In order to estimate its contribution we begin by writing down the amplitude at the quark level:

$$A = \frac{ig}{2\sqrt{2}} \cos \theta_c \bar{u}_d \gamma^\alpha (1 - \gamma^5) u_u \frac{i}{m_W^2} i\sqrt{2} g m_W s_H \frac{i}{m_{\Delta^{++}}^2} \quad (10)$$

$$\times h_{\mu\mu} < leptons > \frac{i}{m_W^2} \frac{ig}{2\sqrt{2}} \sin \theta_c \bar{v}_s \gamma^\alpha (1 - \gamma^5) v_u.$$

We project out the "pseudoscalar" content of A with the operation [10]

$$T_{1b} = C_\pi C_k \sum \bar{u}_u \gamma^5 v_s A \bar{v}_u \gamma^5 u_d \quad (11)$$

where the sum is over the quark spins and C_π and C_k are normalization constants. The amplitude is then evaluated using a simple static constituent quark model. The decay constants are determined by applying the procedure to the decays $K \rightarrow l\nu$ and $\pi \rightarrow l\nu$. The result yields $C_k = if_k m_k / 8m_s m_u$ and $C_\pi = if_\pi m_\pi / 8m_u m_d$. The method can be tested by applying it to the amplitude of Fig. 1(a) and calculating the decay rate for that amplitude alone. In that case the result is the same as that found from the earlier calculation for the amplitude given in Eq. (9).

Applying this procedure to Eq. (10) yields

$$T_{1b} = -1.2 \times 10^3 G_F^{3/2} C_\pi C_k m_\pi^2 m_k E_\pi < leptons > \frac{s_H h_{\mu\mu}}{m_{\Delta^{++}}^2}. \quad (12)$$

The remaining diagrams in Fig. 1 can be evaluated in similar fashion. For example, T_{1e} is given by

$$T_{1e} = i \frac{g^3 h_{\mu\mu}}{2\sqrt{2}} \sin \theta_W \sin \theta_c \cos \theta_c \frac{s_H}{m_W^3 m_{\Delta^+}^2 m_{\Delta^{++}}^2} < leptons >$$

$$\times (m_s - m_u) \bar{v}_s^\alpha (1 + \gamma^5) u_u \bar{u}_d \gamma^\alpha (1 - \gamma^5) v_u (p_{++} + p_-)^\alpha. \quad (13)$$

where p_{++} is the momentum of the incoming Δ^{++} and p_- the momentum of the outgoing Δ_- .

It is clear that with reasonable values for the Higgs masses the contributions from these graphs will be considerably smaller than that from the first two. Therefore we calculate the rate using only the diagrams in Fig 1(a,b). After squaring the amplitude and doing the spin sum we integrate over phase space as follows:

$$R = \frac{1.1 \times 10^{-18}}{4(2\pi)^3} \frac{1}{m_k^3} \left(\frac{s_H h_{\mu\mu}}{m_{\Delta^{++}}^2} \right)^2 \int ds \lambda^{1/2}(s, m_k^2, m_\pi^2) \lambda^{1/2}(s, m_{\mu\mu}^2, m_{\mu\mu}^2) E_\pi^2 p_1 \cdot p_2. \quad (14)$$

Here $s = (p_1 + p_2)^2$, p_1 and p_2 are the muon momenta, and as usual $\lambda(x, y, z) = x^2 + y^2 + z^2 - 2xy - 2xz - 2yz$. Evaluation of the integral yields the following result for the rate:

$$R = 6 \times 10^{-2} \left(\frac{s_H h_{\mu\mu}}{m_{\Delta^{++}}^2} \right)^2 \text{GeV}^4 \frac{1}{\text{sec}}. \quad (15)$$

III. RESULTS AND DISCUSSION

It is possible to develop models that incorporate Higgs triplets where the $\Delta^{++}W^-W^-$ coupling is allowed only with an enabling non-zero vacuum expectation value for the neutral member [7]. In that case one may obtain a limit for h_{ee} through the Majorana mass term in neutrinoless nuclear double beta decay, similar to that shown in Eq. (3). The assumption that the vev is zero avoids problems with maintaining $\rho = 1$, but in that case graphs like Figs. 1(a,b) would not be allowed. However, couplings of the triplet to vector bosons of the type shown in Eq. (6) are a natural consequence of the model being discussed here.

At the same time, the coupling of Higgs to fermions is determined by the overlap of the mass-eigenstate Higgs fields with the doublet, and the H_5 does not have such overlap. The H_3 does and that makes graphs in Figs. 1(c-h) possible, but the contributions to the rate are smaller by at least a factor $\sim (m_k/m_{\Delta^+})^4$.

It is clear then that the rate will be dominated by Figs. 1(a,b), with the result shown in Eq. (15) and with a branching ratio given by

$$BR = 0.9 \times 10^{-9} \left(\frac{s_H h_{\mu\mu}}{m_{\Delta^{++}}^2} \right)^2 \text{GeV}^4. \quad (16)$$

The existing experimental limit [11] for this branching ratio is 1.5×10^{-4} , corresponding to

$$s_H h_{\mu\mu} \leq 1.5 \times 10^2 (m_{\Delta^{++}}/\text{GeV})^2. \quad (17)$$

This result is not invalidated by the previous limits in Eqs. (3,4,5), but it is not exactly a giant step forward either. Taking the values $m_{\Delta^{++}} \sim 100 \text{ GeV}$ [7] and $s_H h_{\mu\mu} \sim 5 \times 10^{-2} m_{\Delta^{++}}$ (near to and not excluded by the limits of Eqs. (4,5)) yields a branching ratio of 2×10^{-16} ,

which is discouragingly small but of the same order of magnitude as the rate for kaon double beta decay induced by light or heavy Majorana neutrinos and reported in earlier publications [4] [5].

In conclusion, we have calculated the rate for $K^\pm \rightarrow \pi^\mp \mu^\pm \mu^\pm$ that can be expected for a decay mediated by doubly-charged Higgs. The branching ratio for reasonable values of the model's parameters is extremely small compared to the experimental limit, but of the same order of magnitude as that induced by virtual Majorana neutrinos. Since the question of lepton-number violation is still outstanding, these processes warrant further theoretical and experimental study.

This work was supported in part by the Natural Sciences and Engineering Research Council of Canada.

-
- [1] D. Smith, C. Picciotto and D. Bryman, Phys. Rev. D **8**, 342 (1973).
 - [2] R. N. Mohapatra and J. D. Vergados, Phys. Rev. D **24**, 1713 (1981); J. Schechter and J. W. F. Valle, Phys. Rev. D **25**, 2951 (1982).
 - [3] L. Wolfenstein, Phys. Rev. D **26**, 2507 (1982).
 - [4] John N. Ng and A. N. Kamal, Phys. Rev. D **18**, 3412 (1978); A. Halprin, P. Minkowski, H. Primakoff and S. P. Rosen, Phys. Rev. D **13**, 2576 (1976).
 - [5] J. Abad, J. G. Esteve and A. F. Pacheco, Phys. rev. D **30**, 1488 (1984).
 - [6] J. F. Gunion, R. Vega, and J. Wudka, Phys. Rev. D **42**, 1673 (1990).
 - [7] J. F. Gunion, Proceedings of the Santa Cruz Workshop on e^-e^- Physics at the NLC, Santa Cruz (1995).
 - [8] J. F. Gunion, J. Grifols, A. Mendez, B. Kayser and F. Olness, Phys. Rev. D **40**, 1546 (1989).
 - [9] M. L. Swartz, Phys. Rev. D **40**, 1521 (1989).

[10] S. Richardson and C. Picciotto, Phys. Rev. D **52**, 6342 (1995); T. Goldman and W. J. Wilson, Phys. Rev. D **15**, 709 (1977).

[11] Particle Data Group, R. M. Barnett *et al.*, Phys. Rev. D **54**, 1 (1996).

Figure Captions

FIG. 1. Diagrams contributing to $K^+ \rightarrow \pi^- \mu^+ \mu^+$.

